



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Let $b^2 \cos^2 C - b^2 + c^2 = 1^2$. Then $c^2 = 1^2 + b^2 \sin^2 C$. Assume $1 = x^2 - y^2$, and $b \sin C = 2xy$. Then $a = \sqrt{(b^2 - 4x^2 y^2) \pm (x^2 - y^2)}$.

Let $b^2 - 4x^2 y^2 = s^2$, $b + 2xy = ks$, and $b - 2xy = s/k$. Then $b = s(k^2 + 1)/2k$, $xy = s(k^2 - 1)/4k$.

Taking $s = 4mk$ we obtain the following formulae:

$$b = 2m(k^2 + 1), \quad c = x^2 + m^2(k^2 - 1)^2/x^2, \\ a = 4mk \pm [x^2 - m^2(k^2 - 1)^2/x^2], \quad C = \cos^{-1}[2k/(k^2 + 1)].$$

As an example, assume $x = 3$, $m = 3$, $k = 2$. We find $a = 24$, $b = 30$, $c = 18$, $\cos C = \frac{4}{5}$.

II. Solution by G. B. M. ZERE, A. M., Ph. D., Parsons, W. Va.

$m^2 + n^2$, $m^2 - n^2$, $2mn$ are the sides of any right triangle. Let $m = pn$.

$\therefore n^2(p^2 + 1)$, $n^2(p^2 - 1)$, $2n^2 p$ are the sides.

The equation for the general value of θ is

$$n^2(p^2 - 1)\cos\theta + 2n^2 p \sin\theta = n^2(p^2 + 1).$$

Let $\sin\beta = \frac{n^2(p^2 - 1)}{n^2(p^2 + 1)}$, $\cos\beta = \frac{2n^2 p}{n^2(p^2 + 1)}$, then the equation becomes

$$\cos\beta \cos\theta + \sin\beta \sin\theta = 1, \quad \cos(\theta - \beta) = 1.$$

$$\therefore \theta - \beta = 2\pi n_1, \quad \theta = 2\pi n_1 + \beta.$$

As θ must be less than $\frac{1}{2}\pi$, the general value of θ is $\theta = \beta$.

Also solved by J. B. Gregg, M. Sc., C. E., Senecaville, Ohio.

PROBLEMS FOR SOLUTION.

ALGEBRA.

192. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College Defiance, O.

What is the difference between the squares of the two *infinite* continued fractions $\left(3 + \frac{1}{6 + \text{etc.}}\right)$ and $\left(2 + \frac{1}{4 + \text{etc.}}\right)$?

193. Proposed by SAUL EPSTEIN, Ph. D., Chicago, Ill.

Professor Goursat states (*Transactions of the American Mathematical Society*, January, 1904, p. 111) that if a_1, a_2, \dots, a_n ; h_1, h_2, \dots, h_n are two sequences, the h 's being all positive, then $\sum \frac{a_i^2}{h_i^2} \geq \frac{(\sum a_i)^2}{\sum h_i}$. Prove this.

194. Proposed by L. E. DICKSON, Ph. D., The University of Chicago.

In the determination of the canonical forms of Abelian transformations modulo p , one is led to the type $[b_1, b_2, b_3]$:

$$\begin{aligned}\xi_1' &= \xi_1, \quad \eta_1' = b_1\xi_1 + \eta_1 + b_2\xi_2 + \eta_2 + b_3\xi_3 + \eta_3, \quad \xi_2' = \xi_2 - \xi_1, \\ \eta_2' &= \eta_2 + b_2\xi_2, \quad \xi_3' = \xi_3 - \xi_1, \quad \eta_3' = \eta_3 + b_3\xi_3.\end{aligned}$$

Find its period and determine the conditions under which it is conjugate with $[c_1, c_2, c_3]$ under Abelian transformation.

GEOMETRY.

219. Proposed by H. F. MacNEISH, A.B., Assistant in Mathematics, University High School, Chicago, Ill.

Draw a line through a given point which shall divide a given quadrilateral into two equivalent parts; (1) when the point lies in a side of the quadrilateral, (2) when the point is without, (3) within the quadrilateral.

220. Proposed by G. B.M. ZERR, A. M., Ph. D., Parsons, West Va.

Two triangles are circumscribed to a given triangle ABC , having their sides perpendicular to the sides of the given triangle. Prove that the two triangles are equal, and find the area of these triangles.

DIOPHANTINE ANALYSIS.

120. Proposed by L. E. DICKSON, Ph. D., Assistant Professor of Mathematics, The University of Chicago.

Find the prime numbers p for which $x^2 - pxz - px - z + p^2 - 3 = 0$ has more than two sets of positive integral solutions x, z , each $< p$.

AVERAGE AND PROBABILITY.

152. Proposed by L. C. WALKER, A.M., Professor of Mathematics, Colorado School of Mines, Golden, Col.

A square hole is cut through the center of a sphere of radius r . Show that the average volume removed is $\frac{1}{12}r^3(23\sqrt{2}-28)$.

MISCELLANEOUS.

143. Proposed by R. L. MOORE, A. M., Fellow in Mathematics, The University of Chicago.

Does there exist a function, of single valued inverse, satisfying the following conditions: (1) if $u=f(v)$, $w=f(u)$ then $w=f(v)$; (2) given $u_1=f(u)$, $v_1=f(v)$, $w_1=f(w)$; if the triple w, v, w can be regarded as the lengths of the sides of a triangle, the triple u_1, v_1, w_1 cannot be similarly regarded.

144. Proposed by IRA M. DeLONG, Professor of Mathematics in the University of Colorado, Boulder, Col.

Determine the number of distinct spherical polygons of n sides formed by arcs of n given great circles on a sphere, each arc to be greater than 0 degrees, and less than 360 degrees, and no two sides of any polygon to lie on the same circle.